Images Videos Maps News Shopping Gmail more -

Sian in



"immobilized antibody" "FLAG pepl 🕱

instant is on v

About 10,900 results (0.12 seconds)

Advanced search

Everything

Images

Videos

News

Shopping

More

Alexandria, VA Change location

Show search tools

Scholarly articles for "immobilized antibody" "FLAG peptide" linker probe



... using a streptavidin sensor surface seifassembled - Pavlickova - Cited by 45
Directed evolution of high-affinity antibody mimics - . . . - Xu - Cited by 154

A streptavidin-biotin-based microarray platform for ... - Pavlickova - Cited by 8

A Streptavidin-Biotin-Based Microarray Platform for Immunoassays

by P Pavlickova - Cifed by 8 - Related articles acid), to which biotin is covalently coupled via linker molecules [N-hydroxy- ... Dissolve the biotinylated peptide M42 probe in PBST buffer to obtain a series of concen The detection limit for the immobilized antibody (Ab27, Ab28) probes Einhauer, A. and Jungbauer, A. (2001) The FLAG peptide, a versatile ...

www.springerlink.com/index/tw0r5437434046h4.pdf -Similar

(PDF) Research Report

File Format: PDF/Adobe Acrobat - Quick View by P Pavlickova - 2003 - Cited by 45 - Related articles Oct 25, 2002 ... lently coupled via linker molecules with Immobilized Antibody. Fragments Using the Protein The biotinylated M42 (c = 10 µg/mL) probe was immobilized onto FLAG peptide, a versatile fusion tag for the ... www.biotechniques.com/multimedia/archive/.../03341m93 10366/mmobilize your own antibody on

CMC Activity Center: Proteomics: Approaches: CMC-Nature Gateway

The sequence binds to an immobilized antibody directed against the FLAG sequence The FLAG peptide, a versatile fusion tag for the purification of ...

www.cellmigration.org/resource/proteomics/prot_approaches.shtml - Smilar

Rapid and sensitive identification of epitopecontaining peptides ...

by CS Raska - 2003 - Cited by 16 - Related articles doi:10.1016/S1044-0305(03)00405-7 | How to Cite or Link Using DOI containing an immobilized antibody against the c-myc epitope (EQKLISEEDL). the Use of Bioreactive Mass Spectrometer Probe Tips in Protein Characterization. ... A. Einhauer and A. Jungbauer, The FLAG Peptide, a Versatile Fusion Tag for the ... linkinghub.elsevier.com/retrieve/pii/S1044030503004057

Generic method for quantification of FLAG-tagged fusion proteins ...

by C Mersich - 2007 - Cited by 7 - Related articles doi:10.1016/j.jbbm.2007.01.004 | How to Cite or Link Using DOI ...

Imkinghub.etsevier.com/retrieve/pii/S0165022X07000048 -Similar

Show more results from elsevier.com

Aces

Peptide/Oligo Conference Progression from Discovery through Preclinical & Clinical Development www.ibclifesciences.com/TIDES

Streptavidin Mag. Beads 4X Higher Binding, 2 for 1 Offer... Less Expensive. Free Papers www.solulink.com

Antibody Immobilization Build Your Own Assays For Vacuum-Based Separation. www.bio-rad.com

THE™ Flag Tag Antibody Flag Tag Monoclonal Antibody The Best Quality in Industry www.genscript.com

DYKDDDDK Tag Antibodies High quality and low cost. 35 years experience. www.iollab.com

Antibody Immobilization sòlid support, starts from \$900 www.cellmosaic.com

See your ad here »

<u>Labeling Substance and Chimera Substance</u>, <u>Process for Preparing</u>...

by K. Shinya - 2005

Dec 13, 2007 ... A method for labeling a **probe** substance, wherein the **probe** substance beads with an **immobilized antibody** specifically recognizing the epitope A **linker** Y (refer to FIG. 3) used to extend the **Flag peptide** (symbol ... www freepatentsonline.com/y2007/0287152.html - Cached

Design and Characterization of Stimuli-Responsive FLAG-tag ...

by C Pöhner - 2006 - Cifed by 3 - Related articles Sep 5, 2008 ... The amount of **immobilized antibody** was determined via the sensor readout (in RU), the interaction of the **FLAG peptide** with the monoclonal antibody 4E11. Photochem. Photobiol. 1994, 59, 491– 496. Direct **Link**: ...

onlinelibrary wiley.com > ... > Journal Home > Vol 22 Issue 4 > Abstract

PLoS ONE: Adherent Monomer-Misfolded SOD1

by Y Watanaba - 2008 - Cited by 3 - Related articles Since the first report of a link between the SOD1 mutations and FALS [3], After SDS electrophoresis, proteins were also **probed** for using The immunoprecipitated samples were eluted with 3×**FLAG peptide** (Sigma). ... **Immobilized antibody**/bound antigen was collected by centrifugation (5000 g for 1 min). ...

www.plosone.org/.../info%3Adoi%2F10.1371% 2Fjournal.pone.0003497 - Cached

immunoenzymometric Assay for a Small Molecule,11-Deoxycortisol...

by N Kobayashi- Block all acs.org results - 2006 - Cifed by 13 - Related articles

Feb 28, 2006... captured by both an **immobilized antibody** and an enzyme-labeled antibody; ... connecting the VH and VL domains via a flexible **linker** peptide. The sequences encoding the His 6 tag,25 the **FLAG peptide**,26 and ... pubs.acs.org/doi/abs/10.1021/ac061866f

Hybridoma and monoclonal antibody for use in an immunoaffinity ...

... hybridoma cell line designated 4E11, which is specific for DYKDDDDK flag peptide, is employed. The sequence of the linker appears below: ##STR1## ...

www.patentgenius.com/patent/5011912.html - Cached

1 2 3 4 5 6 7 8 9 10 Next

Search Help Give us feedback

Google Home - Advertising Programs - Business Solutions - Privacy - About Google